arXiv:1408.3243v1 [math.NT] 14 Aug 2014

JACKSON'S INTEGRAL OF MULTIPLE HURWITZ-LERCH ZETA FUNCTIONS

SU HU AND MIN-SOO KIM

Abstract. Using the Jackson integral, we obtain the q-integral analogue of the Raabe type formula for the multiple Hurwitz-Lerch zeta functions. Our result generalize q-integral analogue of the Raabe type formulas for the Hurwitz zeta functions in [N. Kurokawa, K. Mimachi, and M. Wakayama, Jackson's integral of the Hurwitz zeta function, Rend. Circ. Mat. Palermo (2) 56 (2007), no. 1, 43–56]. During the proof we also obtain a new formula on the relationship between the higher and lower orders Hurwitz zeta functions.

1. Introduction

Raabe's 1843 formula for Gamma function is as follows:

$$\int_0^1 \log\Big(\frac{\Gamma(x+t)}{\sqrt{2\pi}}\Big)dt = x\log x - x, \quad x \geq 0$$

(see [8, p. 89]).

The Hurwitz zeta function $\zeta(s,a)$ is defined by

$$\zeta(s,a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s} \quad (\mathrm{Re}(s) > 1;\ a > 0). \tag{1.1}$$

It also satisfies the Raabe type formula:

$$\int_0^1 \zeta(s, x+t)dt = \frac{x^{1-s}}{s-1}.$$

Using the Volkenborn integral, Cohen and Friedman [3] proved the p-adic Raabe formulas for the Diamond log Gamma function and Washington-Hurwitz zeta function.

The Jackson integral is, when $q > 1$, defined as

$$\int_0^1 f(a)d_q a = (q-1)\sum_{n=1}^{\infty} f(q^{-n})q^{-n} \tag{1.2}$$

2000 *Mathematics Subject Classification.* 11M36.

Key words and phrases. Jackson integral, Raabe type formula, multiple Hurwitz-Lerch zeta function.

(see [5]).

Using the Jackson integral, Kurokawa, Mimachi and Wakayama [7] proved q-integral analogue of the Raabe type formulas for the Hurwitz zeta functions and log Gamma functions.

The multiple Hurwitz-Lerch zeta function $\Phi_k(z,s,a)$ is defined by

$$\Phi_k(z,s,a)=\sum_{m_1,\ldots,m_k=0}^{\infty}\frac{z^{m_1+\cdots+m_k}}{(m_1+\cdots+m_k+a)^s} \tag{1.3}$$

for

$$a\in\mathbb{C}\setminus\mathbb{Z}_0^-;\quad s\in\mathbb{C}\text{ when }|z|<1;\quad \mathrm{Re}(s)>k\ (k\in\mathbb{N})\text{ when }|z|=1.$$

Here $\mathbb{Z}_0^-=\{0,-1,-2,\ldots\}$, $\mathbb{N}$ is the set of positive integers, and $\mathbb{C}$ is the set of complex numbers (see [2]).

Letting $k=1$ in (1.3), we obtain the Hurwitz-Lerch zeta function $\Phi_1(z,s,a)=\Phi(z,s,a)$. Furthermore, when $k=1$ and $z=1$, (1.3) also yields the Hurwitz zeta function $\Phi_1(1,s,a)=\zeta(s,a)$. In addition, letting $k=1, z=1$ and $a=1$ in (1.3), the Riemann zeta function is obtained (for details, see [1, 4, 9, 10]).

In [6], by expressing the sums of products of the Apostol-Bernoulli polynomials in terms of the special values of multiple Hurwitz-Lerch zeta functions at nonpositive integers, we obtained an analogy of Euler's sums of products identity for Apostol-Bernoulli numbers.

In this note, by using the Jackson integral, we extend Kurokawa, Mimachi and Wakayama's q-integral analogue of the Raabe type formula for the Hurwitz zeta functions to the multiple case.

The multiple Lipschitz-Lerch zeta functions $\mathrm{Li}_k(z,s)$ will be involved in our formula, which is defined as

$$\mathrm{Li}_k(z,s)=\sum_{m_1,\ldots,m_k=1}^{\infty}\frac{z^{m_1+\cdots+m_k}}{(m_1+\cdots+m_k)^s}, \tag{1.4}$$

where $s\in\mathbb{C}$ when $|z|<1$, $\mathrm{Re}(s)>k$ $(k\in\mathbb{N})$ when $|z|=1$.

Letting $k=1$ and $z=1$ in (1.4), the Riemann zeta function is obtained.

Denote by $[n]_q=\frac{q^n-1}{q-1}$. Our main result is as follows:

Theorem 1.1. *Let k be an positive integer, $0\le r<k$, and $\Phi_k(z,s,a)$ be the multiple Hurwitz-Lerch zeta function. Then*

$$\int_0^1\Phi_k(z,s,a)d_qa=\frac{1}{[1-s]_q}+\sum_{r=0}^{k-1}\binom{k}{r}\sum_{l=0}^{\infty}\binom{-s}{l}\frac{\mathrm{Li}_{k-r}(z,s+l)}{[l+1]_q} \tag{1.5}$$

is established in the region $a\in\mathbb{C}\setminus\mathbb{Z}_0^-;|z|<1$ and $\mathrm{Re}(s)<1$.

And

$$\int_0^1 \Phi_k(z,s,k-a)d_qa = z^{-k}\sum_{l=0}^{\infty}(-1)^l\binom{-s}{l}\frac{\mathrm{Li}_k(z,s+l)}{[l+1]_q}, \tag{1.6}$$

$$\int_0^1 \Phi_k(z,s,k+a)d_qa = z^{-k}\sum_{l=0}^{\infty}(-1)^l\binom{-s}{l}\frac{\mathrm{Li}_k(z,s+l)}{[l+1]_q} \tag{1.7}$$

are established in the region $a \in \mathbb{C}\setminus\mathbb{Z}_0^-; s \in \mathbb{C}$ *when* $|z|<1$; $\mathrm{Re}(s) > k$ $(k \in \mathbb{N})$ *when* $|z|=1$.

Letting $k=1$ and $z=1$ in Theorem 1.1, we recover the following Kurokawa, Mimachi and Wakayama's results in [7].

Corollary 1.2 (Kurokawa, Mimachi and Wakayama)**.** *Let*

$$\zeta(s,a) = \sum_{n=0}^{\infty}(n+a)^{-s}$$

be the Hurwitz zeta function, analytically continued to all $s \in \mathbb{C}$ *from the region* $\mathrm{Re(s)} > 1$. *Then we have*

$$\int_0^1 \zeta(s,a)d_qa = \frac{1}{[1-s]_q} + \sum_{l=0}^{\infty}\binom{-s}{l}\frac{\zeta(s+l)}{[l+1]_q} \tag{1.8}$$

for $\mathrm{Re(s)} < 1$, *and*

$$\int_0^1 \zeta(s,1-a)d_qa = \sum_{l=0}^{\infty}(-1)^l\binom{-s}{l}\frac{\zeta(s+l)}{[l+1]_q} \tag{1.9}$$

for all $s \in \mathbb{C}\setminus\{1\}$.

During the proof Theorem 1.1, we also obtain the following result on the relationship between the higher and lower orders Hurwitz zeta functions.

Theorem 1.3. *Let* $\Phi_k(z,s,a)$ *be the multiple Hurwitz-Lerch zeta function. Then*

$$\Phi_k(z,s,a) = \frac{1}{a^s} + \sum_{r=0}^{k-1}\binom{k}{r}z^{k-r}\Phi_{k-r}(z,s,a+(k-r)) \tag{1.10}$$

is established in the region $a \in \mathbb{C}\setminus\mathbb{Z}_0^-; s \in \mathbb{C}$ *when* $|z|<1$; $\mathrm{Re}(s) > k$ $(k \in \mathbb{N})$ *when* $|z|=1$.

2. Proof of Theorem 1.3

When $k=3$, we have the following decomposition:

$$\begin{aligned}
\Phi_3(z,s,a) &= \sum_{m_1,m_2,m_3=0}^{\infty} \frac{z^{m_1+m_2+m_3}}{(m_1+m_2+m_3+a)^s} \\
&= \sum_{m_1=1}^{\infty}\sum_{m_2,m_3=0}^{\infty} \frac{z^{m_1+m_2+m_3}}{(m_1+m_2+m_3+a)^s} \\
&\quad + \sum_{m_2,m_3=0}^{\infty} \frac{z^{m_2+m_3}}{(m_2+m_3+a)^s} \\
&= \sum_{m_1,m_2=1}^{\infty}\sum_{m_3=0}^{\infty} \frac{z^{m_1+m_2+m_3}}{(m_1+m_2+m_3+a)^s} \\
&\quad + \sum_{m_1=1}^{\infty}\sum_{m_3=0}^{\infty} \frac{z^{m_1+m_3}}{(m_1+m_3+a)^s} \\
&\quad + \sum_{m_2,m_3=0}^{\infty} \frac{z^{m_2+m_3}}{(m_2+m_3+a)^s} \\
&= \sum_{m_1,m_2,m_3=1}^{\infty} \frac{z^{m_1+m_2+m_3}}{(m_1+m_2+m_3+a)^s} \\
&\quad + \sum_{m_1,m_2=1}^{\infty} \frac{z^{m_1+m_2}}{(m_1+m_2+a)^s} \\
&\quad + \sum_{m_1,m_3=1}^{\infty} \frac{z^{m_1+m_3}}{(m_1+m_3+a)^s} \\
&\quad + \sum_{m_1=1}^{\infty} \frac{z^{m_1}}{(m_1+a)^s} \\
&\quad + \sum_{m_2,m_3=0}^{\infty} \frac{z^{m_2+m_3}}{(m_2+m_3+a)^s}.
\end{aligned} \tag{2.1}$$

Notice that

$$\begin{aligned}
\sum_{m_2,m_3=0}^{\infty} \frac{z^{m_2+m_3}}{(m_2+m_3+a)^s} &= \sum_{m_2=1}^{\infty}\sum_{m_3=0}^{\infty} \frac{z^{m_2+m_3}}{(m_2+m_3+a)^s} \\
&\quad + \sum_{m_3=0}^{\infty} \frac{z^{m_3}}{(m_3+a)^s},
\end{aligned} \tag{2.2}$$

so we have

$$\begin{aligned}\sum_{m_2,m_3=0}^{\infty}\frac{z^{m_2+m_3}}{(m_2+m_3+a)^s} &= \sum_{m_2,m_3=1}^{\infty}\frac{z^{m_2+m_3}}{(m_2+m_3+a)^s}\\ &\quad+\sum_{m_2=1}^{\infty}\frac{z^{m_2}}{(m_2+a)^s}\\ &\quad+\sum_{m_3=1}^{\infty}\frac{z^{m_3}}{(m_3+a)^s}+\frac{1}{a^s}.\end{aligned}\tag{2.3}$$

Combining (2.1) and (2.3), we obtain

$$\begin{aligned}\Phi_3(z,s,a) &= \frac{1}{a^s}+\binom{3}{2}\sum_{m_2=1}^{\infty}\frac{z^{m_2}}{(m_2+a)^s}\\ &\quad+\binom{3}{1}\sum_{m_2,m_3=1}^{\infty}\frac{z^{m_2+m_3}}{(m_2+m_3+a)^s}\\ &\quad+\binom{3}{0}\sum_{m_1,m_2,m_3=1}^{\infty}\frac{z^{m_1+m_2+m_3}}{(m_1+m_2+m_3+a)^s}.\end{aligned}\tag{2.4}$$

In what following, we use the above reasoning (the symmetry of the multiple sum $\sum_{m_1,\ldots,m_k=0}$ in the m_j's variables) to prove an analogue of (2.4) for general k.

Denote by $S=\{m_1,\ldots,m_k\}$. Then

$$\begin{aligned}\Phi_k(z,s,a) &= \sum_{m_1,\ldots,m_k=0}^{\infty}\frac{z^{m_1+\cdots+m_k}}{(m_1+\cdots+m_k+a)^s}\\ &= \frac{1}{a^s}+\sum_{r=0}^{k-1}\sum_{\{m_{i_1},\ldots,m_{i_{k-r}}\}\subset S}\ \sum_{m_{i_1},\ldots,m_{i_{k-r}}=1}^{\infty}\frac{z^{m_{i_1}+\cdots+m_{i_{k-r}}}}{(m_{i_1}+\cdots+m_{i_{k-r}}+a)^s}.\end{aligned}\tag{2.5}$$

Since

$$\sum_{m_{i_1},\ldots,m_{i_{k-r}}=1}^{\infty}\frac{z^{m_{i_1}+\cdots+m_{i_{k-r}}}}{(m_{i_1}+\cdots+m_{i_{k-r}}+a)^s} = \sum_{m_{r+1},\ldots,m_k=1}^{\infty}\frac{z^{m_{r+1}+\cdots+m_k}}{(m_{r+1}+\cdots+m_k+a)^s},$$

from (2.5), we have

$$\Phi_k(z,s,a) = \frac{1}{a^s}+\sum_{r=0}^{k-1}\binom{k}{r}\sum_{m_{r+1},\ldots,m_k=1}^{\infty}\frac{z^{m_{r+1}+\cdots+m_k}}{(m_{r+1}+\cdots+m_k+a)^s}.\tag{2.6}$$

From (2.6), we get

(2.7)

$$\begin{aligned}\Phi_k(z,s,a) &= \frac{1}{a^s}+\sum_{r=0}^{k-1}\binom{k}{r}z^{k-r}\\ &\quad\times\sum_{m_{r+1},\ldots,m_k=1}^{\infty}\frac{z^{(m_{r+1}-1)+\cdots+(m_k-1)}}{((m_{r+1}-1)+\cdots+(m_k-1)+a+(k-r))^s}\\ &= \frac{1}{a^s}+\sum_{r=0}^{k-1}\binom{k}{r}z^{k-r}\Phi_{k-r}(z,s,a+(k-r)).\end{aligned}$$

This completes the proof of Theorem 1.3.

3. Proof of Theorem 1.1

To prove Theorem 1.1, we need the following lemma.

Lemma 3.1. *Let k be an positive integer, $0\le r<k$, and $\Phi_k(z,s,a)$ be the multiple Hurwitz-Lerch zeta function. Then*

$$\int_0^1\Phi_{k-r}(z,s,a+(k-r))d_qa = z^{-(k-r)}\sum_{l=0}^{\infty}\binom{-s}{l}\frac{\mathrm{Li}_{k-r}(z,s+l)}{[l+1]_q}$$

is established in the region $a\in\mathbb{C}\setminus\mathbb{Z}_0^-; s\in\mathbb{C}$ when $|z|<1$; $\mathrm{Re}(s)>k$ $(k\in\mathbb{N})$ when $|z|=1$.

Proof. Using (1.2) and (1.4), we obtain the following equality

$$\begin{aligned}
&\int_0^1 \Phi_{k-r}(z,s,a+(k-r))d_qa \\
&= (q-1)\sum_{n=1}^{\infty} \Phi_{k-r}(z,s,q^{-n}+(k-r))q^{-n} \\
&= (q-1)z^{-(k-r)}\sum_{n=1}^{\infty}\sum_{m_{r+1},\dots,m_k=1}^{\infty} \frac{z^{m_{r+1}+\cdots+m_k}}{(m_{r+1}+\cdots+m_k+q^{-n})^s}q^{-n} \\
&= (q-1)z^{-(k-r)}\sum_{n=1}^{\infty}\sum_{m_{r+1},\dots,m_k=1}^{\infty} \frac{z^{m_{r+1}+\cdots+m_k}}{(m_{r+1}+\cdots+m_k)^s} \\
&\quad\times\left(1+\frac{q^{-n}}{m_{r+1}+\cdots+m_k}\right)^{-s} q^{-n} \\
&= z^{-(k-r)}\sum_{n=1}^{\infty}\sum_{m_{r+1},\dots,m_k=1}^{\infty} \frac{z^{m_{r+1}+\cdots+m_k}}{(m_{r+1}+\cdots+m_k)^s} \\
&\quad\times\sum_{l=0}^{\infty}\binom{-s}{l}\left(\frac{1}{m_{r+1}+\cdots+m_k}\right)^{l} q^{-n(l+1)}(q-1) \\
&= z^{-(k-r)}\sum_{l=0}^{\infty}\binom{-s}{l}\frac{\mathrm{Li}_{k-r}(z,s+l)}{[l+1]_q}.
\end{aligned}$$

This completes the proof of Lemma 3.1. □

Now we prove (1.5). Suppose $a \in \mathbb{C}\setminus\mathbb{Z}_0^-$; $|z|<1$ and $\mathrm{Re}(s)<1$. From Theorem 1.3, we have

(3.1)
$$\begin{aligned}
\int_0^1 \Phi_k(z,s,a)d_qa &= \int_0^1 \frac{1}{a^s}d_qa \\
&\quad+\sum_{r=0}^{k-1}\binom{k}{r}z^{k-r}\int_0^1 \Phi_{k-r}(z,s,a+(k-r))d_qa.
\end{aligned}$$

Thus from the well-known formula (for example, see [7]),

$$\int_0^1 \frac{1}{a^s}d_qa = \frac{1}{[1-s]_q},$$

and Lemma 3.1, we obtain (1.5).

Now we prove (1.6). Suppose $a \in \mathbb{C}\setminus\mathbb{Z}_0^-; s\in\mathbb{C}$ when $|z|<1$; $\mathrm{Re}(s)>k$ $(k\in\mathbb{N})$ when $|z|=1$. As in the prove of Lemma 3.1, we

have

$$\begin{aligned}
&\int_0^1 \Phi_k(z,s,k-a)d_qa \\
&= (q-1)\sum_{n=1}^{\infty}\Phi_k(z,s,k-q^{-n})q^{-n} \\
&= (q-1)\sum_{n=1}^{\infty}\sum_{m_1,\dots,m_k=0}^{\infty}\frac{z^{m_1+\cdots+m_k}}{(m_1+\cdots+m_k+k-q^{-n})^s}q^{-n} \\
&= (q-1)z^{-k}\sum_{n=1}^{\infty}\sum_{m_1,\dots,m_k=1}^{\infty}\frac{z^{m_1+\cdots+m_k}}{(m_1+\cdots+m_k-q^{-n})^s}q^{-n} \\
&= (q-1)z^{-k}\sum_{n=1}^{\infty}\sum_{m_1,\dots,m_k=1}^{\infty}\frac{z^{m_1+\cdots+m_k}}{(m_1+\cdots+m_k)^s} \\
&\quad\times\left(1-\frac{q^{-n}}{m_1+\cdots+m_k}\right)^{-s}q^{-n} \\
&= z^{-k}\sum_{n=1}^{\infty}\sum_{m_1,\dots,m_k=1}^{\infty}\frac{z^{m_1+\cdots+m_k}}{(m_1+\cdots+m_k)^s} \\
&\quad\times\sum_{l=0}^{\infty}(-1)^l\binom{-s}{l}\left(\frac{1}{m_1+\cdots+m_k}\right)^l q^{-n(l+1)}(q-1) \\
&= z^{-k}\sum_{l=0}^{\infty}(-1)^l\binom{-s}{l}\frac{\mathrm{Li}_k(z,s+l)}{[l+1]_q},
\end{aligned}$$

which completes the proof of (1.6). The same procedure proves (1.7).

References

[1] T. M. Apostol, *On the Lerch zeta function*, Pac. J. Math. **1** (1951), 161–167.

[2] J. Choi, D. S. Jang, and H. M. Srivastava, *A generalization of the Hurwitz-Lerch zeta function*, Integral Transforms Spec. Funct. **19** (2008), no. 1-2, 65–79.

[3] H. Cohen, E. Friedman *Raabe's formula for p-adic gamma and zeta functions*, Ann. Inst. Fourier (Grenoble) **58** (2008), no. 1, 363–376.

[4] K. Dilcher, *Sums of products of Bernoulli numbers*, J. Number Theory **60** (1996), 23–41.

[5] F.H. Jackson,*On q-definite integrals*, Quart. J. 41, 193-203.

[6] M.-S. Kim and S. Hu, *Sums of products of Apostol-Bernoulli numbers*, Ramanujan J. **28** (2012), no. 1, 113–123.

[7] N. Kurokawa, K. Mimachi, and M. Wakayama, *Jackson's integral of the Hurwitz zeta function*, Rend. Circ. Mat. Palermo (2) **56** (2007), no. 1, 43–56.

[8] N. Nielsen, Handbuch der Theorie der Gammafunktion, Chelsea, New York, 1965, reprint of 1906 edition.

[9] H. M. Srivastava and J. Choi, *Zeta and q-Zeta Functions and Associated Series and Integrals*, Elsevier Science Publishers, Amsterdam, London and New York, 2012.

[10] K. S. Williams and N. Y. Zhang, *Special values of the Lerch zeta function and the evaluation of certain integrals*, Proc. Amer. Math. Soc. **119** (1993), 35–49.

Department of Mathematics and Statistics, McGill University, 805 Sherbrooke St. West, Montréal, Québec, H3A 2K6, Canada

E-mail address: hus04@mails.tsinghua.edu.cn, hu@math.mcgill.ca

Division of Cultural Education, Kyungnam University, 7(Woryeong-dong) kyungnamdaehak-ro, Masanhappo-gu, Changwon-si, Gyeongsangnam-do 631-701, South Korea

E-mail address: mskim@kyungnam.ac.kr